

# Rhodora

JOURNAL OF THE  
NEW ENGLAND BOTANICAL CLUB.

Conducted and published for the Club, by

BENJAMIN LINCOLN ROBINSON, Editor-in-chief.

FRANK SHIPLEY COLLINS

MERRITT LYNDON FERNALD } Associate Editors.

HOLLIS WEBSTER

WILLIAM PENN RICH

EDWARD LOTHROP RAND } Publication Committee.

Vol. 17.

December, 1915.

No. 204.

## CONTENTS

Range Extension of <i>Ceanothus sanguineus</i> . <i>O. A. Farwell</i> .	229
Characters and Range of <i>Carex laevivaginata</i> . <i>M. L. Fernald</i>	231
<i>Arenaria stricta</i> in the White Mountains. <i>A. S. Pease</i> .	232
<i>Botrychium virginianum</i> v. <i>europaeum</i> in America. <i>M. L. Fernald &amp; H. St. John</i>	233
<i>Bidens connata</i> petiolata. <i>R. W. Woodward</i> . . . .	234
Dr. Gates's Mutation Factor in Evolution (Review). <i>E. M. East</i>	235
Errata . . . . .	238
Index . . . . .	239

Boston, Mass.

1052 Exchange Building.



Providence, R. I.

Preston and Rounds Co.

**RHODORA.**—A monthly journal of botany, devoted primarily to the flora of New England. Price, \$1.50 per year (domestic and foreign); single copies 15 cents. Prices of Volumes 1 and 2 (1899 and 1900) on application. All remittances by check or draft, except on Boston or New York, must include ten cents additional for cost of collection. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be gladly received and published to the extent that the limited space of the journal permits. Forms will be closed five weeks in advance of publication. Authors (of more than one page of print) will receive 25 copies of the issue in which their contributions appear. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to

B. L. ROBINSON, 3 Clement Circle, Cambridge, Mass.

Subscriptions, advertisements, and business communications to

W. P. RICH, 300 Massachusetts Avenue, Boston, Mass.

Single copies may be had from

E. L. RAND, Corresponding Sec'y N. E. Botanical Club,

1052 Exchange Building, Boston, Mass.

Entered at Boston, Mass., Post office as Second Class Mail Matter

---

**KEY TO NEW ENGLAND TREES**, Wild and Commonly Cultivated, based primarily upon leaf characters, by J. FRANKLIN COLLINS and HOWARD W. PRESTON. Price 40c. net. PRESTON & ROUNDS Co., Providence, R. I.

---

**CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS, 1885 TO DATE.**

For American taxonomists and all students of American plants the most important supplement to the Index Kewensis, this catalogue in several ways exceeds the latter work in detail, since it lists not merely the flowering plants, but pteridophytes and cellular cryptogams, and includes not merely genera and species, but likewise subspecies, varieties and forms. A work of reference invaluable for larger herbaria, leading libraries, academies of sciences, and other centers of botanical activity. Issued quarterly, at \$15.00 per 1000 cards.

GRAY HERBARIUM of Harvard University,

Cambridge, Mass., U. S. A.

---

**CHECK LIST OF GRAY'S MANUAL, 7th EDITION**, compiled by M. A. DAY. Leatherette. Pocket size. Invaluable for collector's memoranda and herbarium records. Published and sold by the GRAY HERBARIUM, Cambridge, Mass. Price postpaid 20 cts. each. Ten copies \$1.50.

---

**BOTANICAL BOOKS.** The undersigned will sell cheap, or exchange for Books on BIRDS or INSECTS his present supply of Books treating of Plants, including complete sets of "Rhodora" and Contributions from U. S. National Herbarium.

JOHN D. SHERMAN, JR.,

403 Seneca Avenue, Mt. Vernon, N. Y.

---

*Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. by 3-4 in. 1 year \$4.00, 6 months \$2.50.*

# Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

---

Vol. 17.

December, 1915.

No. 204.

---

## RANGE EXTENSION OF CEANOTHUS SANGUINEUS.

OLIVER A. FARWELL.

IN the Synoptical Flora and in Piper's Flora of the State of Washington the distribution of *Ceanothus sanguineus*, Pursh is given as from Brit. Columbia to N. California and Idaho. Howell gives the same range but extends it eastward to Montana. Rydberg, in the Flora of Montana, says it occurs only on the western slope of the main range of the Rocky Mountains. Coulter's Manual, 1st. Ed., credits it to the region of the Missouri and its tributaries but it is not given a place in the 2nd. edition. Pursh, who first described it from material collected by Lewis, gives it as "Near the Rocky Mountains, on the banks of the Missouri"; in so far as this statement by itself is concerned, it might mean either the eastern or western slope of the Rockies or both; I have not access to any records that might determine the point in question. The plant, however, has been considered to belong exclusively to the northwestern region west of the main range of the Rocky Mountains. Its discovery, therefore, in the Keweenaw Peninsula, in Michigan, is of more than local interest. I first collected it in fruit in August, 1886, near Copper Harbor. At that time, being young in years and botanical experience, my main object was to make each species I found agree with some one of those enumerated in Gray's Manual; so it naturally found a resting place in the species cover of *C. Americanus* where it remained forgotten until 1914. Having in that year had occasion to examine critically my material of the eastern species I at once observed that the Keweenaw plant was not *C. Americanus*. An investigation convinced me that it was either *C. sanguineus* or a new species. On a trip to the Lake Superior region



early in July of this year (1915) I found the shrub in full bloom. It proved to be *C. sanguineus*. I examined a section of territory about half a mile long by as much wide and found the plant to be quite plentiful; the probabilities are that it may be found over a much larger area. It is found in rocky woods, the rock being of trap rock formation; the woods are mostly of evergreen trees but have a good sprinkling of oak, birches, willows, poplars, maples and other deciduous trees and shrubs. It is, of course, indigenous to this region. The possibilities of its having been an introduction from west of the Rockies are so remote as to be negligible. Plants from the eastern slope of the Rockies might be introduced to the Lake Superior region by way of the rivers and streams which find an outlet through Lake Winnipeg, Winnipeg River, the Lake of the Woods, Rainy Lake River, and Pigeon River to Lake Superior but this line of travel would not transport the seed of a plant from the western slope, nor would the Missouri carry the seed from that region to Lake Superior. The same remarks apply to another plant, the *Mimulus moschatus*, Dougl., which is native to the same regions; it may have been introduced further east as an escape from cultivation but the argument will not apply here, since, so far as I have been able to learn, it was never cultivated in the Copper district of Michigan unless gathered for the purpose from the local native plant. The only probable explanation of such widely separated stations is that the species in preglacial times were more generally distributed but that the ice of the glacial period destroyed intermediate stations.

DEPARTMENT OF BOTANY,

PARKE, DAVIS & COMPANY,

DETROIT, MICHIGAN.

## THE CHARACTERS AND RANGE OF *CAREX* *LAEVIVAGINATA*.

M. L. FERNALD.

IN 1909, in his treatment of *Carex* in Engler's *Pflanzenreich*, Küken-thal described from Biltmore, North Carolina, *Carex stipata*, var. *laevivaginata*.<sup>1</sup> In typical *C. stipata* the broadish flaccid leaves have the scarious cross-puckered sheath slightly prolonged at the orifice into a very thin membrane; but in var. *laevivaginata*, as described by Küken-thal, the leaves are firmer and narrower (3 mm. broad), with the sheath plane (not cross puckered) and thickened at the orifice; and the inflorescence is shorter than in typical *C. stipata*. Subsequently, but without amplification of the characters, Mackenzie has elevated the plant to specific rank as *C. laevivaginata* (Kükenth.) Mackenzie<sup>2</sup> and has extended the range to Maryland.

That *Carex laevivaginata* is a distinct species of definite range now seems fairly clear, for a considerable mass of specimens in the Gray Herbarium and the Herbarium of the New England Botanical Club shows it to have not only the characters already emphasized but some others to which attention is here directed, and to occur somewhat generally in damp rich woods and swamps of the Alleghanian region. The most striking character of the plant, besides its firmer and narrower leaves and smooth sheaths with firm often cartilaginous orifice, is its much greener and simpler inflorescences. In *C. stipata* the crowded often somewhat paniculate inflorescence has a yellowish color and in age becomes quite brown; and the perigynia are about 4 mm. long. In *C. laevivaginata*, on the other hand, the inflorescence is simpler and looser, at first pale-green, but in maturity with a very slight brownish tinge; and the perigynia are 5–6.5 mm. long, thus giving the cylindric head an even more muricate appearance than in *C. stipata*.

*Carex stipata* is a plant of the Canadian and Transition Zones, occurring in wet habitats from Newfoundland to British Columbia, southward to the District of Columbia, Virginia, the Great Lake states, Iowa, Kansas, New Mexico and California; with an extremely large

<sup>1</sup> Kükenth. in Engler, *Pflanzenr.* iv. Fam. 20, 172 (1909).

<sup>2</sup> Mackenzie in Britton & Brown, *Ill. Fl.* ed. 2, i. 371 (1913).



variety, var. *uberior* C. Mohr, in the Louisianian area from South Carolina to Florida and Texas. *C. laevivaginata*, on the other hand, is essentially Alleghanian, occurring from Massachusetts to North Carolina, Tennessee, and Illinois. Judged by its representation in herbaria it is much less common in New England than *C. stipata*. The herbarium material immediately accessible shows 88 sheets of *C. stipata* from New England, but only 11 sheets of *C. laevivaginata* representing the following 6 stations. MASSACHUSETTS: Purgatory Swamp, Norwood, June 16, 1878, *Faxon*, June 21, 1896, *E. F. Williams*, June 27, 1897, *Sidney Harris*; Williamstown, May 30, 1898, *J. R. Churchill*, June 17, 1901, *E. F. Williams*; swamp, Stockbridge, June 28, 1900, *Ralph Hoffmann*. RHODE ISLAND: without definite locality, *Olney*; Cat Swamp, Providence, July 2, 1893, *J. F. Collins*. CONNECTICUT: North Cromwell or Farmington, 1878, *Chas. Wright* (both *C. stipata* and *C. laevivaginata* with one label). Judge Churchill has recently informed the writer that *C. laevivaginata* is also found in rich soil near the Merrimac at Bradford, Massachusetts. The species is, however, so little known as yet that further information as to its local distribution and its habitat is needed.

#### GRAY HERBARIUM.

ARENARIA STRICTA IN THE WHITE MOUNTAINS.—In the seventh edition of Gray's Manual *Arenaria stricta* Michx. is reported as found on Mt. Washington, N. H., the citation being based upon a specimen collected by Mr. W. H. Manning. That this reference, however, is due to a transposition of labels and that the specimen upon which it is founded really came from Glens Falls, N. Y., has been convincingly shown (RHODORA, xi, 184-5) by Professor Fernald, who adds that the 'summit of a hill, Holderness, N. H.,' where the plant was collected in July, 1891, by Dr. R. C. Manning, is, so far as he knows, the only New Hampshire station for this species. It was of interest to me, therefore, on 19 July, 1915, while I was collecting on Hart's Ledge, at the southern extremity of the Montalban Range in the lower part of Crawford Notch in Coös and Carroll counties, thirteen miles south of Mt. Washington and about twenty-five miles northeast of Holderness, to find an abundance of this plant growing in woods, on damp rocks, along a steep and somewhat intermittent watercourse. This station is on the southeast side of the ledge in the township of

Bartlett, Carroll County. A more extended visit on 9 September of the same year disclosed the plant on similar damp rocks on the west slope and also in the coarse gravels of the summit (2009 feet), both places being a little west of the county line in Hadley's Grant, Coös County. At the present, then, we may probably regard this ledge as the northeastern limit of the species. Near *A. stricta* in the Carroll County station was its congener, *A. groenlandica* (Retz.) Spreng., which is not uncommon on the mountains of the Montalban Range, but which is perhaps seldom found in so incongruous a situation as here, growing under the shade of red oak trees! Other species of interest noted upon the ledge were *Panicum latifolium* L., *Oryzopsis racemosa* (Sm.) Ricker, *Polygonum scandens* L., *P. Douglasii* Greene (in great abundance on both dry and damp ground, in the sun and in the shade), a slender native *Chenopodium* matching the description of *C. leptophyllum* Nutt., var. *oblongifolium* Wats., *Cardamine parviflora* L., *Geranium Robertianum* L., *G. Bicknellii* Britton, *Arctostaphylos Uva-ursi* (L.) Spreng., and *Specularia perfoliata* (L.) A. DC. Of these the two *Polygonums*, the *Chenopodium*, and the *Arctostaphylos* are new to Coös County, while the *Panicum*, the *Cardamine*, and the *Specularia* are unreported from that county but on Hart's Ledge miss inclusion within its limits by only a few hundred feet. Specimens of all the plants mentioned are to be deposited in the herbarium of the New England Botanical Club. It may be remarked that the vicinity of Bartlett, where plants of the warmer district of east central New Hampshire mingle with those of the colder areas of Crawford Notch and the Montalban and Rocky Branch Ranges, and where the numerous lower mountains are diversified by frequent cliffs and extensive open ledges, offers many attractions to the collector.—ARTHUR STANLEY PEASE, Urbana, Illinois.

THE OCCURRENCE OF *BOTRYCHIUM VIRGINIANUM*, var. *EUROPAEUM* IN AMERICA.—About the Gulf of St. Lawrence, especially on the west coast of Newfoundland, the south coast of the Labrador Peninsula, and on the Gaspé Peninsula of Quebec, there occurs a grape-fern clearly related to *Botrychium virginianum* (L.) Sw., but differing in the much less dissected segments of the sterile frond and, in its most characteristic development, in its heavy or firm texture. The plant proves to be a very close match for the European material passing



as *B. virginianum*, a plant which, as represented in the Gray Herbarium and as shown by the plates illustrating the Scandinavian and Russian specimens, departs from the common Alleghanian and eastern Asiatic material in exactly the points indicated above. This European plant was set off by Presl as *B. anthemoides*,<sup>1</sup> but has subsequently been treated as *B. virginianum*, var. *europaeum* Ångström.<sup>2</sup>

Besides the material from the region of the Gulf of St. Lawrence, characteristic specimens have been seen from the Thunder Bay District, Ontario, and from northern New England. In New England some of the plants seem to be exactly intermediate, as judged by the cutting of the frond, between *B. virginianum* and var. *europaeum*, but south of this area all the plants seem to be typical *B. virginianum*. The plant with the less dissected frond, being the only representative of the species in Europe and occurring in North America chiefly north of the range of typical *B. virginianum*, seems to the writers to be well separated as a variety which should be called *B. virginianum* (L.) Sw., var. *europaeum* Ångström.—M. L. FERNALD and HAROLD ST. JOHN.

*BIDENS CONNATA PETIOLATA*.—At Franklin, Connecticut, in low meadows, *B. connata petiolata* occurs with noticeably large heads and with achenes considerably exceeding the length of 4–6 mm. given for the species in the last edition of the Manual. In the Franklin plants the mature central achenes are 8–9 mm. long. Very few central achenes shorter than 8 mm. were found in the many heads examined by the writer. The measurements refer to large, well developed heads. The awns, too, are longer than in the species, as well as stouter, and the two pairs are often of equal length. Short golden-yellow rays are usually developed early but soon fall away. The plants are abundant in Franklin Meadows, and very uniform in habit. They show well a difference between variety and species, which seems, so far as the writer's observation extends, to be constant, but is, however, more readily recognized in the field than in the herbarium. In the typical form of the species the head is low and hemispherical, while in the variety it is taller and cylindrical, or at full maturity slightly broadened upward. In fresh plants the contrast is striking. Specimens have been deposited in the Gray Herbarium.—R. W. WOODWARD, New Haven, Connecticut.

<sup>1</sup> Presl, Abh. böhm. Ges. ser. 5, v. 323 (1848).

<sup>2</sup> Ångström, Botaniska Notiser (1854) 68.



THE MUTATION FACTOR IN EVOLUTION: WITH PARTICULAR REFERENCE TO *OENOTHERA*. By R. RUGGLES GATES.<sup>1</sup> Since DeVries discovered the two aberrant forms of *Oenothera Lamarckiana* that he called *brevistylis* and *laevifolia* at Hilversum in 1887, the genus *Oenothera* probably has been studied more intensively than any other similar group of plants. As evidence of this analytic activity a literature of some five hundred titles has been produced. Every botanist appreciates the numerous facts that have been brought to light regarding the classification, distribution, gross morphology, cytology and genetics of the genus, and owing to the volume of the publications everyone is thankful when compilations are made. DeVries has brought together the greater part of these facts in *Die Mutations-theorie* and *Gruppenweise Artbildung*. The volume under consideration supplements these two works by a fairly complete consideration of the taxonomy, cultural history and cytology of the group. In addition the author brings together the data from his own extensive pedigree culture, and urges, rather didactically at times, the conclusions he has drawn from them.

The book purports to be a reconnaissance of the theory of Evolution by Mutation, it really is wholly a consideration of the biology of the *Oenotheras*. The author rests his case on the one group of facts. DeVries did not make this mistake; for, as Cervantes says, "It is the part of a wise man to keep himself today for tomorrow, and not to venture all his eggs in one basket." DeVries did indeed lay great stress upon his work with the evening primroses, but he did not overlook numerous other props for his hypothesis,—props so sturdy that in the opinion of some, the *Oenothera* investigations might be disregarded without weakening the edifice. Perhaps few biologists adhere strictly to DeVries' views of Evolution. In particular it might be mentioned that he did not go far enough in distinguishing between germinal and somatic variations, and that he has not kept pace with the facts regarding inheritance. At the same time, it must be admitted that in addition to the great stimulus to experimental biology that his work effected, DeVries made two great general contributions. He showed the frequency with which germinal changes of comparatively great size occur, and why they are not swamped by intercrossing. But these generalizations make no new Evolution theory. They merely extend and modify Darwin's ideas insofar as these new facts tend to change the emphasis the latter placed upon particular types of variation.

These changes in viewpoint may be made with total disregard for the *Oenothera* work. In fact, perhaps few angiosperm genera could have been selected which are so fundamentally unsuited for genetic work from which broad conclusions are to be drawn as *Oenothera*. As Gates shows, numerous aberrant types of chromosome distribution occur at gametogenesis. Presumably many of the daughter cells

<sup>1</sup> London. Macmillan, 1915. pp. I-XIV + 353. Figs. 114.

formed by such irregular division are aborted. Thus there is a selective elimination of potential gametes. Further, many of the pollen grains, and possibly the egg cells, are not functional. Again, an extremely large percentage—often the majority—of the zygotes formed are not viable. This will be apparent to anyone who takes the trouble to make germination tests of *Oenothera* seeds. Even after discarding the many seeds that casual examination shows to be worthless, it is seldom that over sixty percent of the remainder produce mature plants, and the germination may drop as low as five percent. Now, it is quite likely that the only useful laws of heredity will be those which like the laws of physics and chemistry are mathematical descriptions of cycles of events from which predictions of what must occur under like circumstances may be made. Is it strange then, that many biologists are cautious when asked to accept as a basis for such descriptions, breeding results from plants like the *Oenotheras* where only a small portion of the facts can be known owing to the immense number of potential plants lost through the abortion of both zygotes and gametes? It is like asking a chemist to accept theories as to the structural formulae of organic compounds upon which only determinations of nitrogen and oxygen have been made.

These facts are boldly disregarded by the author in his chapters on "Hybridisation and Hereditary Behavior" and "The Relation between Hybridisation and Mutation." The *Oenotheras*, he says, have four main types of hereditary behavior: "(1) mutation crosses, (2) Mendelian splitting, (3) blending and modification of characters, and (4) twin hybrids." Much of the discussion under these heads is a severe arraignment of Mendelism, but the author's contempt for the Mendelian theory of heredity is not that bred of familiarity. If the reviewer has not misconceived matters, the author's idea of Mendelian segregation is enticingly simple. If when two organisms are crossed, The  $F_1$  generation is uniform and the  $F_2$  generation comprises two types in the ratio of three to one, the inheritance is Mendelian. No circumstance whatever may modify the definiteness of these phenomena under pain of their disqualification as examples of Mendelian inheritance.

It would be rash to assert that Mendelism even in the broadened sense with which the word is used today, covers all types of inheritance. Correns' and Baur's experiments on chromatophores and Goldschmidt's work on the gypsy moth indicate the possibility of inheritance through the cytoplasm, while Mendelian data parallel chromosome distribution.

On the other hand, it is even more bold to assert that inheritance in *Oenothera* is definitely nonmendelian. They cannot be placed in either category with certainty, but it ought to be emphasized that no single fact discovered by those who have made pedigree cultures of the group, precludes a Mendelian interpretation. Gates' arguments against Mendelian interpretation of heredity in the evening primroses



reduce to four: (1) It has been definitely proven that *O. Lamarchiana* is not a hybrid; (2) Constant ratios of the usual Mendelian type are not found; (3) Splitting often occurs in the  $F_1$  instead of the  $F_2$  generations, sometimes with the production of two types unlike the parents (twin hybrids); and, (4) Chromosome differences appear.

In regard to the first point, it may be said that the finding of a single herbarium specimen hardly constitutes proof that the plant in question is or is not a hybrid. The second point is covered by our remarks on the selective elimination of zygotes and gametes. The third argument overlooks a fundamental tenet of Mendelism. Crosses split first in the second hybrid generation, solely when the eggs on the one hand and the sperms on the other hand entering into the so-called  $F_1$  generation were of the same factorial constitution. The mere act of producing a cross does not necessarily make the next generation an  $F_1$  generation. The fact that *O. Lamarchiana* gives off variants is to many an indication that a cross between it and another form does not produce an  $F_1$  hybrid. The frequency or infrequency with which the aberrant types appear proves nothing as long as we are ignorant of the potentialities of the non-functional gametes and zygotes. With this ignorance to contend with, the variability of the ratios is even an argument in favor of segregation rather than mutation, for it must be remembered that the same types continually reappear, whereas by DeVriesian theory mutations are equally likely to occur in any direction. The fourth argument of the author is really a question of terminology. If the behavior of the chromosomes is the efficient cause of Mendelian phenomena, then even aberrant mitoses at reduction are in the broad sense Mendelian. They hold possibilities of variation without that true germinal change which may be pictured as a chemical reconstitution independent of the mechanics of cell division.

In calling attention to these points, however, the reviewer does not wish to have it understood that he denies mutation. Wide variations and narrow variations as opposed to mere adaptive fluctuations certainly appear, and some of these variants may have been produced independently of "slips" in cell division. But it seems to him unwise to make the case rest upon the *Oenothera* data. The author speaks truly when he says that "biology has passed the stage when single evolutionary factors, no matter how insistently urged or how brilliantly advocated, can be held accountable for the great diversity of life which we see around us." — E. M. EAST.

## ERRATA.

- Page 30, line 3; for Hamden read Hampden.  
" 31, " 34; for confirms read confirm.  
" 32, " 1; for Mongaric read Morgani.  
" 66, " 10; for 400 read 700.  
after that insert some of  
" 66, " 11; for between 100 and 200 read of about 400.  
" 66, " 15; after that insert one of  
for station read stations.  
" 66, " 16; for below 600 read of 720.  
" 71, " 16; for northwestern read northeastern.  
" 72, " 12; for TERETOLOGICAL read TERATOLOGICAL.  
" 102, " 20; for heft read Heft.  
" 146, " 29; after L. insert a comma.  
" 148, " 32; after T. & G. insert a comma.  
" 151, " 24; for Brittanica read Britannica.  
" 151, " 25; for Radula read radula.  
" 152, " 38; for Filix-foemina read Filix-femina.  
" 184, " 3; for O. Williamsii read E. Williamsii.  
" 198, " 31; after Gaspé insert and.  
" 219, " 28; for Aehillea read Achillea.

Vol. 17, no. 203, including pages 205 to 228, was issued 7 December, 1915.



# INDEX TO VOLUME 17.

New scientific names are printed in full face type.

- Abies*, 166, 167; *alba*, 62; *americana*, 167; *balsamea*, 60, 65, 152; *canadensis*, 61, 62, 166, 167; *Fraseri*, 60, 165; *pectinata*, 62; *picea*, 62.  
*Acalypha virginica*, 226.  
*Acer carolinianum*, 66; *rubrum*, var. *tridens*, 66.  
*Achillea Millefolium*, 36, 154, 219.  
*Acnida*, 178; *cannabina*, 178, 209, 215, 221; *rhyssocarpa*, 179; *tamariscina*, 179; *tuberculata*, 179.  
*Acrostichum areolatum*, 162.  
*Azelia cassioides*, 134.  
*Agropyron*, 226; *caninum*, var. *tennerum*, 147; *repens*, 146, 147, 213, 226.  
*Agrostis alba*, 147, 150, 209, 212, var. *maritima*, 209, 212, 221, var. *vulgaris*, 153, 209, 212; *hyemalis*, 153, 212.  
*Ailanthus glandulosa*, 217.  
Aizoaceae, 203.  
Albino *Viola rostrata*, 180.  
*Aletris aurea*, 64.  
Algae from the Chincha Islands, Some, 89.  
*Alisma Plantago-aquatica*, 211.  
*Alnus incana*, 150, 152, 153; *mollis*, 148, 152.  
*Alopecurus*, 97.  
Altitudinal Limits in Connecticut, A Correction, 88.  
Amaranth, Bloody, 179.  
Amaranthaceae, 178.  
*Amaranthus*, 179, *blitoides*, 179; *caudatus*, 179; *cruentus*, 179; *gracilis*, 179; *graecizans*, 179; *hybridus*, 179, forma *hypochondriacus*, 179; *Palmeri*, 179; *paniculatus*, 179; *Powellii*, 179; *retroflexus*, 179, 226; *spinosus*, 180; *viridis*, 180.  
*Ambrosia artemisiifolia*, 36, 219, 225, 226; *trifida*, 225; *trifida integrifolia*, 30.  
*Amelanchier laevis*, 152, 153.  
America, The Occurrence, of *Botrychium virginianum*, var. *europeum* in, 233; some new or unrecorded Compositae chiefly of north-eastern, 1.  
American Ostrich Fern, 161; Representatives, *Elymus arenarius* and its, 98; Variations of *Lycopodium annotinum*, 123.  
Amherst, Weeds growing in, 29.  
*Ammophila arundinacea*, 146.  
*Amsonia*, 130.  
*Anagallis arvensis*, 35.  
*Anaphalis margaritacea*, 36, 154, 219.  
*Andromeda acuminata*, 132; *glaucophylla*, 148; *lucida*, 132; *nitida*, 132; *populifolia*, 132; *reticulata*, 132.  
*Anemone parviflora*, 159.  
Annual winter Meeting of the Vermont Botanical Club, 86.  
Anonymos, 130; *aquaticus*, 130; *caroliniensis*, 130, 131, 134; *cassioideis*, 134; *pinnatus*, 131; *repens*, 130; *umbrosus*, 131.  
*Antennaria*, 126; *canadensis*, 125-128, 154, in Pennsylvania, 125; *eucosma*, 159; *fallax*, 127; *neglecta*, 127; *neodioica*, 127; *occidentalis*, 127, 128; *Parlinii*, 127, 128; *petaloidea*, 127; *plantaginifolia*, 36, 37; *spathulata*, 159.  
*Anthemis Cotula*, 36; *tinctoria*, 30.  
*Anthoceros crispulus*, 120.  
*Anthoxanthum odoratum*, 221.  
*Anychia*, 202; *canadensis*, 202; *polygonoides*, 202.  
*Aplectrum hyemale*, 136.  
*Arabis viridis*, 65.  
*Aralia hispida*, 152, 154; *nudicaulis*, 152.  
*Arceuthobium*, 171; *pusillum*, 171.  
*Arctium minus*, 36.  
*Arctostaphylos*, 233; *Uva-ursi*, 233.  
*Arenaria caroliniana*, 66; *groenlandica*, 233; *lateriflora*, 148, 151; *peploides*, var. *robusta*, 146; *stricta*, 232, 233, in the White Mountains, 232.  
*Arethusa*, 150; *bulbosa*, 65, 149; *racemosa*, 136; *spicata*, 136.

- Aristolochiaceae, 171.  
*Arnica alpina*, 19; *chionopappa*, 159; **pulchella**, 18, 19; *tomentosa*, 19.  
*Arrhenatherum elatius*, 146.  
*Artemisia cordata*, 38; *vulgaris*, 219.  
*Asarum*, 171; *canadense*, 171.  
*Asclepias incarnata*, var. *pulchra*, 217.  
*Asparagus officinalis*, 214.  
*Aspidium americanum*, 48; *campylopterum*, 48; *cristatum*, 150; *dilatatum*, 47; *dumetorum*, 47; *intermedium*, 47; *noveboracense*; 152; *spinulosum*, 47, 153, var. *americanum*, 47, 48, var. *dilatatum*, 44, 47, 152, 153, var. *dilatatum*, forma *anadenium*, 45, 48, var. *intermedium*, 152; *Thelypteris*, 150.  
*Asplenium*, 64; *Filix-femina*, 150, 152.  
*Aster*, 14, 17; *acuminatus*, 152; **anticostensis**, 16, 17; *cordifolius*, 36; *divaricatus*, 36; *Douglasii*, 17; *ericoides*, 36, 37; *foliaceus*, 13, 14, 16, 17, var. **arcuans**, 14, 15, var. **crenifolius**, 15, var. *frondeus* 14, 15, var. *Parryi*, 14, 16, var. **subpetiolatus**, 15, 16, var. **subgeminatus**, 16; *Fremonti*, 13; **johannensis**, 13; *juncus*, 13; *lateriflorus*, 36, 218, var. *hirsuticaulis*, 152, 154; *linariifolius*, 36; *longifolius*, 13, 17; *multiflorus*, 36; *nemoralis*, 67-69, var. *Blakei*, 149; *novibelgii*, 13, 36, 217, 218, 221, var. *litoreus*, 147; *paniculatus*, 17, 154; *patens*, 36; *puniceus*, 36, 152, var. **perlongus**, 17; *radula*, 151, 152; *spathulatus*, 17; *spectabilis*, 65; *subulatus*, 218, 221; *tardiflorus*, 16; *undulatus*, 36; *vimineus*, 36, 218.  
*Atriplex*, 83, 85, 176; *arenaria*, 84, 85, 176; *bracteosa*, 176; *dioica*, 84; *halimoides*, 84; *halimus*, 84; *hastata*, 84; *laciniata*, 84, 85; *maritima*, 84, 85; *maritimum*, 84, 85; *mucronata*, 84; new to North America, 83; *patula*, 83, 176, var. *hastata*, 146, 147, 176, 208, 215, 221, var. *littoralis*, 176; *rosea*, 85, 176; *sabulosa*, 84, 85.  
*Avena sativa*, 212.  
*Baccharis halimifolia*, 218, 221.  
Balch, W. E., The field Meeting of the Vermont Botanical and Bird Clubs, 143.  
Balsam, Fir, 165-168.  
*Baptisia tinctoria*, 35.  
*Barbarea vulgaris*, var. *longisiliquosa*, 216.  
Barn yard Grass, 225.  
Bartlett, H. H., Systematic Studies on *Oenothera*,—V, 41.  
*Bassia*, 176; *hirsuta*, 176.  
Beal, W. J., Weeds growing in Amherst, 29.  
Beggars Ticks, 225.  
Bergen, J. Y., Weed Growth and unusual Rainfall, 224.  
Beta, 64.  
*Betula alba*, var. *cordifolia*, 148, 152, var. *papyrifera*, 152; *glandulosa*, 87, var. *rotundifolia*, 87, var. **sibirica**, 87; *nana*, 87,  $\beta$  *sibirica*, 87, var. *sibirica*, 87; *populifolia*, 153, 154, 214; *rotundifolia*, 87.  
*Bidens*, 20, 63; *Beckii*, 63; *bidensoides*, 21; *cernua*, 21, 25, 219, var. *elliptica*, 25, var. *integra*, 25, var. *minima*, 25, var. **oligodonta**, 25; **colpophila**, 21; *comosa*, 23, 24; *connata*, 23, 24, 219, var. **fultior**, 24, var. *petiolata*, 221, 234; *discoidea*, 219; *Eatoni*, 20, 21; *frondosa*, 22, 219, 225, var. **stenodonta**, 22; **heterodoxa**, 23, var. **orthodoxa**, 24; *hyperborea*, 20, 21; in eastern North America, Some anomalous Species and Varieties of, 20; *tripartita*, 22, 23, var. *heterodoxa*, 23.  
Black Grass, 214, 218, 219.  
Blake, S. F., An *Atriplex* new to North America, 83; *Betula glandulosa*, var. *sibirica* (Ledeb.) n. comb., 87; *Moneses uniflora*, var. **reticulata**, 28; A new Form of *Polygala polygama*, 201; Notes on the Genus *Sabatia*, 50; Some neglected Names in Walter's *Flora Caroliniana*, 129.  
Bloody Amaranth, 179.  
*Boehmeria*, 169; *cylindrica*, 169.  
Boston District, Reports on the Flora of the,—XX, 169, XXI, 202.  
Botanical and Bird Clubs, The field Meeting of the Vermont, 143; Club, The 20th annual winter Meeting of the Vermont, 86; Society of Maine, The Josselyn, 128; Society of Maine, The twenty-first annual Meeting of the Josselyn, 203.  
*Botrychium angustisegmentum*, 87, 88, 143; *anthemoides*, 234; *lanceolatum*, 87, 88, 143, in the Yellowstone, 143, var. *angustiseg-*



- mentum, 87; virginianum, 233, 234, var. europaeum, 234, var. europaeum in America, The Occurrence of, 233.
- Brainerd, E., *Viola septentrionalis* in British Columbia, 70.
- Brassica arvensis, 30, 35; Napus, 35.
- British Columbia, *Viola septentrionalis* in, 70.
- Bromus ciliatus, 150; hordeaceus, 30, 38; Kalmii, 71; tectorum, 38.
- Brunnichia cirrhosa, 130.
- Cakile edentula, 35, 146.
- Calamagrostis canadensis, 150.
- Calopogon pulchellus, 149.
- Calypogeia paludosa, 107, 119, 120, forma subaquatica, 120; sphagnicola, 119; tenuis, 119; Trichomanis, 119, 120, var. tenuis, 119.
- Camelina microcarpa, 30.
- Campanula rapunculoides, 36; rotundifolia, 148.
- Cannabis, 169; sativa, 169.
- Capsella Bursa-pastoris, 35, 216.
- Cardamine, 156, 233; hirsuta, subsp. oligosperma, 157, subsp. oligosperma, var. bracteata, 157, subsp. oligosperma, proles lasiocarpa, 157, subsp. oligosperma, proles unijuga, 158; oligosperma, 156, 157, The Varieties of, 156, var. **bracteata**, 157, var. **lucens**, 157, var. **unijuga**, 158; parviflora, 233; unijuga, 156, 158.
- Carex, 204; adusta, 153, 154; brunescens, 152; canescens, var. disjuncta, 148; castanea, 65, 66, 88; communis, 152; canoidea, 150; Crawfordii, 150, var. vigens, 150; crinita, 152, 227, var. gynandra, 150, 152, var. Porteri, 227; debilis, var. Rudgei, 150; diandra, 65; filiformis, 150; from Newfoundland, A new, 158; hormathodes, 150, 214, 221, var. invisus, 214, 221; laevivaginata, 231, 232, Characters and Range of, 231; leptalea, 148; limosa, 148-150; maritima, 147; misandra, 159; **misandroides**, 158, 159; novae-angliae, 65, 152; Oederi, var. pumila, 150, 151; oligosperma, 148; pallescens, 150, 153; pauciflora, 148, 149; pauperula, var. irrigua, 148; pedata, 159; plantaginea, 204; rostrata, var. utriculata, 150; scoparia, 151, 213, var. condensata, 151, var. moniliformis, 151; stellulata, 150, var. angustata, 150, var. cephalantha, 150; sterilis, 148, 150; stipata, 214, 231, 232, var. laevivaginata, 231, var. uberius, 232; tribuloides, 214; trisperma, 148, 152, var. Billingsii, 148, 149; Tuckermanni, 57, 58; var. **niagarensis**, 57-59; Tuckermanni niagarensis, a neglected Sedge, 57; vulpinoidea, 214.
- Carolina Laurel, 50.
- Carpet weed, 226.
- Caryophyllaceae, 37.
- Caulophyllum, 204, 224.
- Ceanothus americanus, 229; sanguineus, 229, 230, Range Extension of, 229.
- Celtis, 169; occidentalis, 169.
- Cephalozia catenulata, 107, 112-116; divaricata, 118; fluitans, 120; macrostachya, 107, 113-117, var. propagulifera, 117; media, 114-116; multiflora, var. elata, 114; reclusa, 112; serriflora, 112, 113; virginiana, 112, 113.
- Cephalozia, 117; Hampeana, 117, var. spinicaulis, 117, forma typica, 117; papillosa, 118; spinicaulis, 107, 117, 118.
- Ceranium clavulatum, 96; minutum, 96; rubrum, 94, 95.
- Cerastium vulgatum, 35, 209, 216.
- Ceratodon purpureus, 210.
- Chaenolobus, 135; undulatus, 135, 137.
- Chaetomorpha Linum, 91.
- Chamaedaphne calyculata, 148.
- Chaptalia semifloscularis, 136, 137; tomentosa, 135-137.
- Characters and Range of Carex laevivaginata, 231.
- Chase, A., A teratological Specimen of Panicum amarulum Hitchc. & Chase, 72.
- Chelidonium majus, 35.
- Chelone glabra, 151.
- Chenopodiaceae, 176.
- Chenopodium, 176, 233; album, 176, 215, 226, var. viride, 176; ambrosioides, 176, var. anthelminticum, 176; Bonus-Henricus, 176; Boscianum, 176; Botrys, 177; capitatum, 177; carinatum, 34; glaucum, 177; graveolens, 177; hybridum, 177; leptophyllum, 177, var. oblongifolium, 233; multifidum, 177; murale, 177; polyspermum, 177; rubrum, 177; urbicum, 177; virgatum, 177.

- Cherokee Plum, 49.  
 Chestnuts, Spanish, 49.  
 Chiloscyphus pallescens, 120.  
 Chinch Islands, Some Algae from the, 89.  
 Chiococca alba, 51.  
 Chiogenes serpyllifolia, 152.  
 Chironia, 129; campanulata, 52; gracilis, 52; lanceolata, 50, 51.  
 Chlora, 55; dodecandra, 55.  
 Chondrus canaliculatus, 92.  
 Chorisporea tenella, 32.  
 Chorizanthe pungens, 175.  
 Chrysanthemum Leucanthemum, var. pinnatifidum, 36, 225.  
 Chrysocoma acaulis, 135.  
 Chrysopsis falcata, 36.  
 Cichorium, 227; Intybus, 38, 220.  
 Cicuta bulbifera, 217; maculata, 221.  
 Circaea, 223; alpina, 152, 223; intermedia, 204; latifolia, 223, and the Asiatic C. quadrisulcata, The Identity of, 222; lutetiana, 222, 223,  $\beta$  canadensis, 223, var. quadrisulcata, 224; maritima, 223; quadrisulcata, 224, The Identity of Circaea latifolia and the Asiatic, 222.  
 Cirsium arvense, 36, 220; discolor, 36, 37; lanceolatum, 220.  
 Cladium mariscoides, 151.  
 Cladophora fascicularis, 91; Hariotina, 91.  
 Clayton's Oaks in the British Museum, Three of, 39.  
 Club, The field Meeting of the Vermont Botanical and Bird, 143, The 20th annual winter Meeting of the Vermont Botanical, 86.  
 Coelopleurum actaeifolium, 146.  
 Collins, F. S., Some Algae from the Chinch Islands, 89; November Flowers, 33.  
 Comandra, 171; livida, 149; umbellata, 171.  
 Common Syringa, 121.  
 Compositae, 1, 37, 222; chiefly of northeastern America. Some new or unrecorded, 1.  
 Conioselinum, 151; chinense, 151.  
 Connecticut, Altitudinal Limits in, a Correction, 88.  
 Conringia orientalis, 30.  
 Convolvulus arvensis, 226; sepium, 151, var. pubescens, 146, 148; speciosus, 131.  
 Coptis trifolia, 152.  
 Cornus canadensis, 146, 152, 154.  
 Correction, Altitudinal Limits in Connecticut, 88; concerning Sagittaria teres Watson and Potamogeton lucens L., 142.  
 Corsinia, 109.  
 Corydalis sempervirens, 35, 154.  
 Couch Grass, 226.  
 Crab Grass, 225.  
 Crataegus, 153, 154.  
 Crepis capillaris, 30.  
 Cruciferae, 37.  
 Cryptogramma Stelleri, 65.  
 Cuscuta Gronovii, 151, 217.  
 Cycloloma, 64.  
 Cyperaceae, 222, 225.  
 Cyperus ferax, 213; strigosus, 225.  
 Cypridium acaule, 152.  
 Cyprus, 50.  
 Cystopteris fragilis, 78.  
 Cytire, 50.  
 Daisy, Ox eye, 225.  
 Dalibarda repens, 152.  
 Danthonia spicata, 153.  
 Datura Tatula, 30.  
 Daucus Carota, 35.  
 Deane, W., A Correction concerning Sagittaria teres Watson and Potamogeton lucens L., 142; Floral Changes in a Salt Marsh during Reclamation, 205; Reports on the Flora of the Boston District,—XX, 169, XXI, 202.  
 Delesseria Hypoglossum, 94.  
 Delphinium carolinianum, 130; consolidata, 35, 38.  
 Dentaria diphylla, 204.  
 Deschampsia flexuosa, 148.  
 Dianthus Armeria, 35.  
 Dicksonia punctilobula, 153.  
 Diervilla Lonicera, 152, 153.  
 Digitalis purpurea, 228.  
 Digitaria humifusa, 225; sanguinalis, 212, 225, 226.  
 Diplophyllum apiculatum, 120.  
 Dipteracanthus ciliolus,  $\gamma$  parviflora, 134.  
 Dirca palustris, 204.  
 Distichlis spicata, 79, 213, 221.  
 Dracoecephalum denticulatum, 134.  
 Drosera longifolia, 149, 150; rotundifolia, 149.  
 Dryas integrifolia, 159.  
 Dryopteris dilatata, 44, 45; Goldiana, 46; intermedia, 44; patens, 64; spinulosa, 44, 48, var. americana, 48, var. dilatata, 44-46, The North American Representatives of, 44, var. glandulosa, 45, var. intermedia, 44.



- Dulichium arundinaceum*, 150, 151.  
 Dwarf Mistletoe, 65.
- East, E. M., The mutation Factor in Evolution; with particular Reference to *Oenothera*, 235.
- Echinochloa crusgalli*, 105, 106, 209, 212, 225; **muricata**, 106, 209, 212; Walteri.
- Ectocarpus Mitchellae*, 91, 92; *virens*, 92.
- Eleocharis acicularis*, 142, 151; *palustris*, 150, 151, var. *vigens*, 150; *tenuis*, 151.
- Elymus ampiculmis*, 99; *arenarius*, 98-103, 146, 147, and its American Representatives, 98, forma *compositus*, 102, var. **compositus**, 102, 103, var. *villosus*, 99-103; *capitatus*, 99, 100; *dives*, 99; *giganteus*, 98, 103; *mexicanus*, 98; *mollis*, 99-101; *sabulosus*, 98; **strigatus**, 102, 103; *villosissimus*, 100.
- Elytraria caroliniensis*, 131, 137, var. **angustifolia**, 131, 137; *virgata*, 131, 137, var. *angustifolia*, 131, 137.
- Emex*, 171; *spinosa*, 171.
- Empetrum nigrum*, 148, 149.
- Endoderma strangulans*, 91.
- Enteromorpha intestinalis*, 91; *prolifera*, 91.
- Epilobium adenocaulon*, 151, 217, var. *perplexans*, 154; *angustifolium*, 154, 217; *coloratum*, 217; *densum*, 151.
- Equisetum arvense*, 150; *scirpoides*, 64; *sylvaticum*, 152.
- Eragrostis capillaris*, 227; Frankii, 227.
- Erechtites hieracifolia*, 209, 218, 219, 225.
- Erigeron canadensis*, 36, 218, 225; *hysopifolius*, 18, var. **villicaulis**, 17; *ramosus*, 36.
- Eriocaulon septangulare*, 68.
- Eriophorum angustifolium*, 149, 151; *callitrix*, 149; *tenellum*, 149; *virginicum*, 149.
- Erodium cicutarium*, 35, 37.
- Errata, 238.
- Eryngium*, 132.
- Eupatorium perfoliatum*, 151; *purpureum*, var. *maculatum*, 151, 218.
- Euphorbia hirsuta*, 30; *humistrata*, 30, 227; *maculata*, 227.
- Euphrasia*, 181-183, 194; *albimontana*, 185; *americana*, 148, 154, 182, 183, 185, 196-198, 200, 201, var. *canadensis*, 183, 195; *arctica*, 182-184, 190, 192, 194, 196; *bottnica*, 187; *canadensis*, 181-184, 195-198, 201; **disjuncta**, 182, 184, 190, 192, 194; *hirtella*, 183, 190, 193; **hudsoniana**, 184, 194, 197; in North America; The Genus, 181; *latifolia*, 181-183, 190, 192-194; *mollis*, 182-184, 190; *nemorosa*, 196, 197, 201; *Oakesii*, 181-183, 185, 186, forma **lilacina**, 185; *officinalis*, 181, 182, 185, 188, 190, 195, 199, 200,  $\gamma$  *alpestris*, b. *arctica*, 193, *officinalis alpina*, 182, var. *hyperborea*, 193,  $\beta$  *latifolia*, 193, var. *mollis*, 190, var. *tatarica*, 182, 185, 193, 198; *orbicularis*, 185; *purpurea*, 182, 184, 186, 189, 190, forma **candida**, 187, var. **Farlowii**, 184, 189, 190, Farlowii, forma *iodantha*, 189, var. **Randii**, 184, 188-190, var. **Randii**, forma **albiflora**, 188; *pusilla*, 182, 185; **Randii**, 148, 183, 188, var. Farlowii, 183, 189; *Rostkoviana*, 183; *salisburgensis*, 195; *stricta*, 182, 184, 196-198, 201, var. **tatarica**, 184, 194, 197, 198; *tatarica*, 193, 198; *Williamsii*, 183-185, var. **vestita**, 184, 186.
- Eusabbatia*, 55.
- Euthamia camporum*, 12; *floribunda*, 12.
- Euthora cristata*, 94.
- Evans, A. W., Notes on New England Hepaticae,—XII, 107.
- Extensions of Range, 227.
- Eyebright, 181, 183.
- Fagopyrum*, 172; *esculentum*, 172; *tataricum*, 172.
- Farwell, O. A., The Hemlock Spruce, 164; Range Extension of *Ceanothus sanguineus*, 229.
- Fellows, D. W., The Josselyn Botanical Society of Maine, 128; The twenty-first annual Meeting of the Josselyn Botanical Society of Maine, 203.
- Fern, The American Ostrich, 161; Ostrich, 161.
- Fernald, M. L., An albino *Viola rostrata*, 180; Altitudinal Limits in Connecticut, a Correction, 88; The American Ostrich Fern, 161; The American Variation of *Lycopodium annotinum*, 123; *Botrychium angustisegmentum* (Pease & Moore), n. comb., 87; The Char-

- acters and Range of *Carex laevivaginata*, 231; Flora of the Vicinity of New York, a Contribution to Plant Geography, 62; The Genus *Euphrasia* in North America, 181; a handy Book on Trees and Shrubs, 103; The Identity of *Circaea latifolia* and the Asiatic *C. quadrisulcata*, 222; Michaux's *Panicum muricatum*, 105; A new *Carex* from Newfoundland, 158; The North American Representatives of *Dryopteris spinulosa*, var. *dilatata*, 44; The Occurrence of *Botrychium virginianum*, var. *europaeum* in America, 233; Some anomalous Species and Varieties of *Bidens* in eastern North America, 20; Some new or unrecorded Compositae chiefly of northeastern America, 1; Two Variations of *Silene antirrhina*, 96.
- Festuca elatior*, 213; *ovina*, 213; *rubra*, 147, 148.
- Field Meeting of the Vermont Botanical Bird Clubs, 143.
- Finger Grass, 225.
- Fir Balsam, 165-168; Silver, 165.
- Fletcher, E. F., Two more foreign Plants found on Wool-waste at Westford, 32.
- Flora of the Boston District, Reports on the, XX, 169, XXI, 202; of the Vicinity of New York, a Contribution to plant Geography, 62.
- Floral changes in a Salt Marsh during Reclamation, 205.
- Flynn, N. F., The 20th annual winter Meeting of the Vermont Botanical Club, 86.
- Fossombronina 108, 109; caespitiformis, 110; cristula, 107, 109-111; echinata, 110; foveolata, 109, 110; Luetzelburgiana, 109.
- Foxtail, Green, 226.
- Frasera, 130.
- Galinsoga parviflora*, 30.
- Galium aparine*, 146, 147; *Claytoni*, 151; *trifidum*, 151.
- Gates, R. R. [Notice of Work], 235.
- Gaultheria procumbens*, 152.
- Gaylussacia baccata*, 148; *dumosa*, var. *Bigeloviana*, 148, 149.
- Gentian, 56.
- Gentiana*, 56; *crinita*, 35.
- Genus *Euphrasia* in North America, 181.
- Geranium Bicknellii*, 233; *pusillum*, 30; *Robertianum*, 35, 233.
- Gerardia cassioides*, 134; *maritima*, 208, 221.
- Gerbera*, 136.
- Geum canadense*, 35.
- Gigartina*, 96; *Chamissoi*, 92; *Lessonii*, 93.
- Glaux maritima*, var. *obtusifolia*, 147.
- Glyceria borealis*, 150; *canadensis*, 72, 150; *grandis*, 150; *laxa*, 150; *nervata*, 150; *pallida*, 150; *septentrionalis*, 213; *Torreyana*, 150.
- Gnaphalium polycephalum*, 36, 219; *uliginosum*, 227; *undulatum*, 135.
- Gomphrena*, 180; *globosa*, 180.
- Gracilaria*, 96; *confervoides*, 93.
- Gramineae, 222; Two Extensions of Range in, 71.
- Grass, Barn yard, 225; Black, 214, 218, 219; Couch, 226; Crab, 225; Finger, 225; Witch, 225.
- Grasses, Range Extensions of two, 159.
- Green Foxtail, 226.
- Grover, F. O., Extensions of Range, 227.
- Habenaria clavellata*, 151; *obtusata*, 152.
- Hamamelis virginica*, 35.
- Handy Book on Trees and Shrubs, 103.
- Hedysarum alpinum*, 159.
- Helianthus grosseserratus*, 225; *laetiflorus*, 219.
- Hemlock, 165, 167, 168; Pines, 50; Spruce, 59-61, 164-168.
- Hepaticae, Notes on New England,—XII, 107.
- Heracleum*, 151; *lanatum*, 151.
- Hexalectris aphylla*, 136; *spicata*, 136.
- Hieracium aurantiacum*, 154; *canadense*, 20, var. *hirtirameum*, 19; *floribundum*, 154.
- Hierochloë odorata*, 146, 147, 212, 221.
- Honey Locust, 49.
- Hordeum jubatum*, 147, 221.
- Horseweed, 225.
- Houstonia caerulea*, 154.
- Howe, I. A., The field Meeting of the Vermont Botanical and Bird Clubs, 143.
- Hubbard, F. T., *Mibora minima* (L.) Desv. at Plymouth, Massachusetts, 97.
- Humulus*, 169; *japonicus*, 169; *Lupulus*, 169.

- Hunnewell, F. W., 2d, *Botrychium lanceolatum* in the Yellowstone, 143.
- Hydrocotyle americana*, 151.
- Hyoseris biflora*, 135.
- Hypericum acutifolium*, 134; *angulosum*, 134; *boreale*, 221; *canadense*, 151; *denticulatum*, 134, 137, var. **acutifolium**, 134, 137, var. **ovalifolium**, 135, 137; *ellipticum*, 151; *virgatum*, 134, 137, var. *acutifolium*, 134, 137, var. *ovalifolium*, 135, 137; *virginicum*, 151.
- Ichnanthus*, 72.
- Identity of *Circaea latifolia* and the Asiatic *C. quadrisulcata*, 222.
- Ilex verticillata*, 150, 152.
- Illecebraceae, 202.
- Illecebrum*, 203; *verticillatum*, 203.
- Impatiens biflora*, 151, 217.
- Insular Variety of *Solidago semper-virens*, 26.
- Ipomoea sagittata*, 131, 137; *spiciosa*, 131, 137.
- Iris Hookeri*, 155; *setosa*, var. *canadensis*, 148; *versicolor*, 151.
- Josselyn Botanical Society of Maine, 128; Botanical Society of Maine, The twenty-first annual Meeting of the, 203.
- Juncoides caroliniae*, 63; *echinatum*, 64; *pilosum*, 63; *saltuense*, 63.
- Juncus*, 214; *acuminatus*, 214; *alpinus*, var. *insignis*, 204; *aristulatus*, 66; *balticus*, var. *littoralis*, 147, 151; *brevicaudatus*, 151; *bufonius*, 147, 151, var. *halophilus*, 147; *conglomeratus*, 64; *effusus*, var. *Pylaei*, 151; *filiformis*, 151; *Gerardi*, 147, 208, 209, 212-216, 218, 220, 221; *Greenei*, 153, 154; *tenuis*, 151, 214; *trifidus*, 65, var. *monanthus*, 65.
- Jungermannia catenulata*, 112, 113; *reclusa*, 112, 113.
- Juniperus communis*, var. *depressa*, 153, 154; *horizontalis*, 148.
- Kalmia angustifolia*, 148; *polifolia*, 148.
- Kantia Trichomanis*, var. *tenuis*, 119.
- Knowlton, C. H., Plants and plant Societies at Roque Bluffs, Maine, 145; Reports on the Flora of the Boston District,—XX, 169, XXI, 202.
- Kobresia caricina*, 159.
- Kochia*, 177; *scoparia*, 177.
- Krigia amplexicaulis*, 135, 137; **bi-flora**, 135, 137.
- Kuhnistera pinnata*, 131.
- Labrador Tea, 149.
- Lachnanthes tinctoria*, 68.
- Lactuca canadensis*, 220; *scariola*, var. *integrata*, 209, 218, 220.
- Lady's Thumb, 225.
- Lapitheia*, 55, 56; *Boykinii*, 54.
- Laportea*, 169; *canadensis*, 169.
- Larch, 65.
- Larix laricina*, 148, 152.
- Lastraea dilatata*, var. *glandulosa*, 45, 47.
- Lathyrus maritimus*, 146; *palustris*, var. *pilosus*, 154.
- Laurel, Carolina, 50.
- Lechea intermedia*, 154.
- Ledum*, 66, 88, 146; *groenlandicum*, 65, 148, 149.
- Leersia oryzoides*, 212.
- Leguminosae, 37.
- Lemna minor*, 214.
- Leontodon*, 146, 227; *autumnalis*, 36, 220, var. *pratensis*, 146.
- Leonurus cardiaca*, 35.
- Lepidium virginicum*, 35, 216.
- Lepiota*, 31, 32; *americana*, 31; *Morgani*, 32; *naucina*, 31; *procera*, 30, 31; *rhacodes*, 31.
- Leucothoe acuminata*, 132, 137; *populifolia*, 132, 137.
- Ligusticum scoticum*, 148.
- Limonium carolinianum*, 35, 147, 217, 221.
- Linaria canadensis*, 36, 209, 218, 220; *vulgaris*, 36, 217.
- Linnæa borealis*, var. *americana*, 152.
- Liriodendron*, 224.
- Listera cordata*, 154.
- Lithospermum carolinianum*, 131, 137; *Gmelini*, 131, 137.
- Locust, Honey, 49.
- Lomaria*, 161.
- Lonicera caerulea*, var. *villosa*, 150; *canadensis*, 152.
- Lophocolea alata*, 107, 111, 112; *bidentata*, 112; *cuspidata*, 111, 112, var. *alata*, 111.
- Lophozia Hatcheri*, 120; *longidens*, 120.
- Loranthaceae, 171.
- Love Lies Bleeding, 179.
- Lupinus perennis*, 71.
- Luzula campestris*, var. *echinata*, 64; *carolinæ*, 63; *pilosa*, 63; *saltuensis*, 63, 152.



- Lychnis dioica*, 216.  
*Lycopodium adpressum*, 66; *alope-curoides*, 66; *annotinum*, 123, 124, 152, The American Variations of, 123, var. **acrifolium**, 124, 125, var. *alpestre*, 124, 125, var. *pungens*, 123-125; *apodum*, 68; *clavatum*, 152, 153; *lucidulum*, 152; *obscurum*, var. *dendroideum*, 152.  
*Lycopus americanus*, 217; *uniflorus*, 151.  
*Lygodium palmatum*, 68.  
*Lyonia lucida*, 132, 137; *nitida*, 132, 137.  
*Lysimachia terrestris*, 151.  
 Macbride, J. F., Range Extensions of two Grasses, 159; *Sisymbrium brachycarpon* and Allies, 138; A white flowered *Spiraea tomentosa*, 142.  
*Macbridea caroliniana*, 132, 137; *pulchra*, 132, 137.  
*Maclura*, 169; *pomifera*, 169.  
*Maianthemum canadense*, 152.  
 Maine, The Josselyn Botanical Society of, 128; Plants and plant Societies at Roque Bluffs, 145; The twenty-first annual Meeting of the Josselyn Botanical Society of, 203.  
*Malva rotundifolia*, 35.  
 Massachusetts, *Mibora minima* (L.) Desv. at Plymouth, 97.  
 Mathews, F. S. [Notice of Work], 103.  
*Matricaria suaveolens*, 146.  
*Matteuccia*, 161, 162; **nodulosa**, 164; *Struthopteris*, 161, 162, 164.  
*Medicago lupulina*, 35.  
*Megalodonta*, 63.  
*Melampyrum lineare*, 149, 152.  
*Melilotus alba*, 35, 217.  
*Mentha arvensis*, 217.  
*Menyanthes*, 150; *trifoliata*, 149, 150.  
*Mertensia maritima*, 146, 147.  
*Mibora*, 97; *minima*, 97, at Plymouth, Massachusetts, 97; *verna*, 97.  
 Michaux, Washington and, 49.  
 Michaux's *Panicum muricatum*, 105.  
*Micranthemum orbiculatum*, 131, 137; **umbrosum**, 131, 137.  
*Microstylis uniflora*, 149.  
*Mimulus moschatus*, 230.  
 Mistletoe, Dwarf, 65.  
*Mitchella repens*, 152.  
 Mock Orange, 121.  
*Mollugo*, 203, 226; *verticillata*, 35, 203, 226, 227.  
*Moneses*, 28; *grandiflora*, 28; *reticulata*, 28, 29; *uniflora*, 28, 29, 152, var. **reticulata**, 28, 29.  
*Monotropa uniflora*, 152.  
*Montia lamprosperma*, 154.  
 Moore, A. H., Two *Philadelphus* Combinations, 121.  
*Morus*, 170; *alba*, 170; *rubra*, 170.  
 Mutation Factor in Evolution; with particular Reference to *Oenothera*, 235.  
 Mycophagist, A rash, 30.  
*Mylia anomala*, 120.  
*Myrica asplenifolia*, 153, 154; *Gale*, 150.  
*Naia flexilis*, 211.  
 Name of the Hemlock Spruce, 59.  
*Nemopanthus mucronata*, 51, 148, 152.  
*Neottia glandulosa*, 136.  
*Nephrodium spinulosum fructuosum*, 45.  
 New Carex from Newfoundland, A, 158.  
 New England Hepaticae, Notes on,—XII, 107.  
 New Form of *Polygala polygama*, 201.  
 Newfoundland, A new Carex from, 158.  
 New York Botanical Garden, The twentieth Anniversary of the, 144.  
*Nitophyllum cryptoneuron*, 95.  
 North America, An *Atriplex* new to, 83; The Genus *Euphrasia* in, 181; *Rumex persicarioides* and its Allies in, 73; Some anomalous species and Varieties of *Bidens* in eastern, 20.  
 North American Representatives of *Dryopteris spinulosa*, var. *dilatata*, 44.  
 Northern *Solidago* in the White Mountains, 72.  
 Notes on New England Hepaticae, —XII, 107; on the Genus *Sabatia*, 50.  
 November Flowers, 33.  
*Nyctaginaceae*, 202.  
*Nymphaea advena*, var. *variegata*, 150.  
 Oak, Red, 30.  
 Occurrence of *Botrychium virginianum*, var. *europaeum* in America, 233.

- Oenothera, 235-237; atrovirens, 41, 42; biennis, 217, 225; **cleistantha**, 41-44; cruciata, 41; Lamarckiana, 235, 237, forma brevistylis, 235, forma laevifolia, 235; muricata, 35, 146, 147, 217; Mutation Factor in Evolution; with particular Reference to, 235; Oakesiana, 41; pumila, 35, 154; **Robinsonii**, 41-44; stenomeris, 41, 42; stenopetala, 41; Systematic Studies on,—V, 41; venosa, 41-44.
- Onoclea, 161; nodulosa, 161, 162, 164; Struthiopteris, 161, 162.
- Orange, Mock, 121.
- Orchis spectabilis, 204.
- Oryzopsis racemosa, 233.
- Osmunda caroliniana, 162; cinnamomea, 150; Claytoniana, 150.
- Ostrich Fern, 161; The American, 161.
- Oxalis Acetosella, 152; corniculata, 154, 226.
- Oxybaphus, 202; floribundus, 202; nyctagineus, 202.
- Ox eye Daisy, 225.
- Paliurus, 49.
- Pallivicia Flotowiana, 120.
- Panicum, 71, 72, 233; amarulum Hitchc. & Chase, A teratological Specimen of, 72; Ashei, 71; barbulatum, 71; boreale, 153; capillare, 212, 225; Commonsianum, 71; crusgalli, var. muricatum, 106; depauperatum, 71; dichotomiflorum, 212; implicatum, 153; latifolium 233; linearifolium, 65; muricatum, 105, 106; pungens, 106; stipitatum, 160; strictum, 71; umbrosum, 71.
- Parentucellia, 194.
- Parietaria, 170; pennsylvanica, 170.
- Paronychia, 203; argyrocoma, var. albimontana, 203.
- Pease, A. S., Arenaria stricta in the White Mountains, 232; A northern Solidago in the White Mountains, 72.
- Pennsylvania, Antennaria canadensis in, 125.
- Pericidium semiflosculare, 135.
- Petalostemum corymbosum, 131, 137; **pinnatum**, 131, 137.
- Petasites, 66, 88; palmata, 66.
- Phalaris arundinacea, 212.
- Phegopteris hexagonoptera, 204; polypodioides, 152.
- Philadelphus acuminatus, 122; Combinations, Two, 121; coronarius, 122, 123, var. **acuminatus**, 122, var. satsumi, 121, var. tomentosus, 122, 123; Gordonianus, 122; grandiflorus, 122; intectus, 123; latifolius, 122, 123; Lewisii, 122; pubescens, 122, var. **intectus**, 123; satumanus, 121, 122; satsumi, 121, 122; tomentosus, 122; Yokohama, 121; Yokohamae, 121.
- Philirea latifolio, 50.
- Phleum alpinum, 78; pratense, 212.
- Phoradendron flavescens, 64.
- Physostegia denticulata, 134, 137; **purpurea**, 134, 137; virginiana, 134.
- Phytolacca, 180; americana, 180; decandra, 34, 180.
- Phytolaccaceae, 180.
- Picea canadensis, 152; glauca, 62; mariana, 148, 149, 171; rubra, 148, 152.
- Pigweed, 226.
- Pilea, 170; pumila, 170.
- Pines, Hemlock, 50.
- Pinus, 166, 167; Abies, 62; alba, 62; balsamea, 59, 60, 165-168; Banksiana, 155; canadensis, 59-62, 166-168; glauca, 62; laxa, 62; Picea, 60.
- Pistatia, 50.
- Plantago decipiens, 63, 147, 148, 218, 221; major, 218; maritima, 63; Rugelii, 227.
- Plants and plant Societies at Roque Bluffs, Maine, 145.
- Pleienta, 55-57; capitata, 54, 55; fasciculata, 55; flexuosa, 55; gentianoides, 55; leucantha, 55; rigida, 55.
- Pleonosporium venustissimum, 95.
- Plocamium, 94, 96; coccineum, 93-95, forma **compactum**, 93, 94, var. flexuosum, 95; leptophyllum, 95, var. flexuosum, 95.
- Pluchea camphorata, 219.
- Plum, Cherokee, 49.
- Plurimaria, 56.
- Plymouth, Massachusetts, Mibora minima (L.) Desv. at, 97.
- Poa annua, 213, 227; compressa, 221; pratensis, 153, 213; triflora, 153, 213.
- Podophyllum, 224.
- Podostemon ceratophyllum, 68.
- Pogonia, 150; ophioglossoides, 149.
- Polygala polygama, A new Form of, 201, var. abortiva, 202, forma **obovata**, 201.

- Polygonaceae, 171, 222.  
 Polygonella, 172; articulata, 34, 172.  
 Polygonum, 172, 233; acre, 172, var. leptostachyum, 172, 215; amphibium, 172, forma Hartwrightii, 172, forma terrestre, 172; arenarium, 172; arifolium, 172; aviculare, 34, 172, 227, var. angustifolium, 172, var. littorale, 173, var. vegetum, 146, 173; Bellardi, 173; Bistorta, 173; Careyi, 173; cili-node, 152, 153, 173; Convolvulus, 173; cuspidatum, 173; Douglasii, 233; dumetorum, 173, forma cristatum, 173; erectum, 173; exsertum, 173, 215, 221; Fowleri, 147; glaucum, 63, 173; Hydro-piper, 173, 215; hydropiperoides, 173, var. strigosum, 173; lapathi-folium, 174, 215, var. incanum, 174, var. nodosum, 174; maritimum, 63, 173; Muhlenbergii, 174; orientale, 174; pennsylvanicum, 174; 215; Persicaria, 34, 174, 215, 225; prolificum, 173, 174, 215, 221; ramosissimum, 174, forma atlanticum, 174, 215, 221; rurivagum, 174; sagittatum, 151, 174, 215; scandens, 174, 233; tenue, 174; tomentosum, 174; virginianum, 174, 224.  
 Polypodium polypodioides, 64; vul-gare, 152.  
 Polysiphonia, 95.  
 Polytrichum commune, 210.  
 Pontederia cordata, 214.  
 Ponthieva glandulosa, 136; **race-mosa**, 136.  
 Populus grandidentata, 214; tremu-loides, 152, 214.  
 Portulaca oleracea, 226.  
 Potamogeton, 66, 211; alpinus, 150; amplifolius, 142; bupleuroides, 210; confervoides, 66; crispus, 210, 211, 221; dimorphus, 211; epihydrus, 150, 210; lucens, 142, A Correction concerning Sagit-taria teres Watson and, 142; natans, 150.  
 Potentilla, 150; argentea, 35; cana-densis, 35, var. simplex, 154; mon-speliensis, 154, 216; nivea, 159; pacifica, 147, 208, 216, 221; palus-tris, var. villosa, 150; recta, 30; tridentata, 148, 154.  
 Pottia truncata, 210.  
 Prasium, 134; incarnatum, 134; pur-pureum, 134.  
 Prenanthes alba, 135; autumnalis, 135, 137; nana, 154; trifoliolata, 152; virgata, 135, 137.  
 Pretz, H. W., Antennaria canadensis in Pennsylvania, 125.  
 Pride of China Plants, 50.  
 Primula farinosa, var. macropoda, 154, 155.  
 Prunella vulgaris, 36, var. lanceolata, 154.  
 Prunus pennsylvanica, 152.  
 Pseudochironia, 55.  
 Pteris aquilina, 153; tremula, 64.  
 Pterocaulon, 135; pycnostachyum, 135, 137.  
 Pterosiphonia dendroidea, 95.  
 Puccinellia angustata, 147; mari-tima, 208, 213, 221.  
 Purslane, 226.  
 Pyrola americana, 152; elliptica, 152; uniflora, 28.  
 Pyrolaceae, 28.  
 Pyrus americana, 152, 153; arbuti-folia, var. atropurpurea, 152; me-lanocarpa, 150; sitchensis, 152, 153.  
 Quercus coccinea, 39; digitata, 39; falcata, 39; Michauxii, 40; mon-tana, 40; Muehlenbergii, 40; Prinos, 40; rubra, 39, 40; velu-tina, 39.  
 Radicula palustris, 216, var. hispida, 216.  
 Ragweed, 225.  
 Rajania, 130; ovata, 130.  
 Ramnus Tree, 50.  
 Range Extension of Ceanothus san-guineus, 229; Extensions of, 227; Extensions of two Grasses, 159.  
 Ranunculus acris, 35; Cymbalaria, 79, 147, 208, 216; pennsylvanicus, 151; repens, 35; sceleratus, 209, 216.  
 Raphanus Raphanistrum, 35.  
 Rash Mycophagist, 30.  
 Red Oak, 39.  
 Rehder, A., The Name of the Hem-lock Spruce, 59.  
 Reports on the Flora of the Boston District,—XX, 169, XXI, 202.  
 Rheum, 174; Rhaponticum, 174.  
 Rhexia Alifanus, 132, 137; glabella, 132, 137.  
 Rhinanthus Crista-galli, var. fallax, 148.  
 Rhododendron canadense, 148.  
 Rhodymenia corallina, 93; flabelli-folia, 93; Palmetta, 94.  
 Rhus typhina, 217.



- Ribes oxyacanthoides*, 152; *prostratum*, 152.  
*Riccia arvensis*, 120.  
*Ricciella crystallina*, 120.  
*Riella*, 109.  
 Roque Bluffs, Maine, Plants and plant Societies at, 145.  
*Rosa nitida*, 150; *rugosa*, 216; *virginiana*, 35, 37, 146-148, 150, 153, 221.  
 Rosaceae, 37.  
*Rubus*, 86, 153; *canadensis*, 153; *Chamaemorus*, 149; *nigricans*, 152, 153; *tardatus*, 152, 153; *triflorus*, 152.  
*Rudbeckia hirta*, 36.  
*Ruellia carolinensis*, 134; *carolinensis*, 134, 137, var. *parviflora*, 134, 137; *ciliosa*, 134, 137, var. *ambigua*, 134, var. *parviflora*, 134, 137.  
*Rumex*, 73, 75, 79, 175, 226; *Acetosa*, 175; *Acetosella*, 175, 215, 226; *altissimus*, 175; *Britannica*, 151, 175; *crispatus*, 77; *crispus*, 175, 215; *fueginus*, 76, 81; *hastatus*, 175; *hispidus*, 150; *idaeus*, var. *aculeatissimus*, 153; *limosus*, 80; *maritimum*, var. *fueginum*, 79; *maritimus*, 73, 75-77, 80, 83, var. ***athrix***, 79, 80, 82, var. *fuegina*, 76, *fueginus*, 77, 79-81, 83, 175; *mexicanus*, 30, 175; *obtusifolius*, 77, 175; *pallidus*, 146, 175; *Patientia*, 146, 151, 175; *persicarioides*, 73-77, 80, 81, 83, 175, and its Allies in North America, 73; *salicifolius*, 79; *verticillatus*, 175.  
*Rynchospora alba*, 148, 151; *capillacea*, var. *leviseta*, 204.  
*Sabatia*, 54-56, 129; *angularis*, 56; *arenicola*, 56; *Boykinii*, 54, 55; *brachiata*, 56; *calycina*, 56; *campanulata*, 52, 56, 57, var. ***grandiflora***, 52, 56; *campestris*, 56; ***capitata***, 54, 55, 57; *carnosa*, 56; *decandra*, 56; *difformis*, 50, 51, 56, 57; *dodecandra*, 54-56; *Elliottii*, 51, 56; *foliosa*, 53, 54, 56; *gentianoides*, 55, 57; *gracilis*, 52; *Harperi*, 53, 54; *lanceolata*, 50, 51; *macrophylla*, 56; *maculata*, 56; Notes on the Genus, 50; ***obtusata***, 54, 56, 57; *Palmeri*, 56; *paniculata*, 51, 56,  $\beta$  *angustifolia*, 51,  $\alpha$  *latifolia*, 51; *Purpusii*, 56; *simulata*, 56; *stellaris*, 56.  
*Sabbatia*, 55; *Boykinii*, 54; *campanulata*, 52; *corymbosa*, 50; *difformis*, 50; *gracilis*, 52, var. *grandiflora*, 52; *grandiflora*, 52; *lanceolata*, 50, 51; *stellaris*, 52.  
*Sagina nodosa*, var. *glandulosa*, 154; *procumbens*, 148.  
*Sagittaria latifolia*, 209, 211, forma *gracilis*, 211, forma *hastata*, 211, forma *obtusata*, 211; *teres*, 142, and *Potamogeton lucens* L., A Correction concerning, 142.  
 St. John, H., *Elymus arenarius* and its American Representatives, 98; An insular Variety of *Solidago sempervirens*, 26; The Occurrence of *Botrychium virginianum* var. *europaeum* in America, 233; *Rumex persicarioides* and its Allies in North America, 73; Some anomalous Species and Varieties of *Bidens* in eastern North America, 20.  
*Salicornia*, 178; *ambigua*, 178; *europaea*, 147, 178, 215, 221, var. *pachystachya*, 178; *mucronata*, 178.  
*Salix alpina*, 136, 137; *arbuscula*  $\alpha$  *erecta*, forma *angustifolia*, 136; *babylonica*, 214; *balsamifera*, 152, 153; *discolor*, 64, 152, var. *eriocephala*, 152; *fusca*, 136; *humilis*, 152; *lanata*, 136; *leirolepis*, 159; *lucida*, 150; *Myrsinites*, 136,  $\beta$  *Jacquiniiana*, 136; *rostrata*, 152; *squamata*, 64; *tristis*, 136, 137.  
*Salsola*, 178; *Kali*, 178, var. *caroliniana*, 178, var. *tenuifolia*, 178.  
 Salt Marsh during Reclamation, Floral Changes in a, 205.  
*Sambucus canadensis*, 218; *racemosa*, 152.  
 Santalaceae, 171.  
*Saponaria officinalis*, 35.  
 Sargent, C. S., Three of Clayton's Oaks in the British Museum, 39; Washington and Michaux, 49.  
*Sarracenia purpurea*, 65.  
*Scheuchzeria*, 149; *palustris*, 148.  
*Schizaea*, 67; *australis*, 68; *filifolia*, 68; *pusilla*, 67, 68.  
*Scirpus americanus*, 147, 213, 221; *atrocinetus*, 151; *campestris*, var. *paludosus*, 208, 213, 221; *debilis*, 63; *georgianus*, 151; *Peckii*, 64; *pedicellatus*, 154; *robustus*, 213, 221; *rubrotinctus*, 151, var. *confertus*, 151; *Smithii*, 63; *validus*, 150.  
*Scleranthus*, 203; *annuus*, 35, 203.  
*Scutellaria caroliniana*, 132, 133; *galericulata*, 151; *integrifolia*, 133;

- ovalifolia, 133; ovata, 132, 133, 137, var. **bracteata**, 134, 137; pilosa, 132, 133; versicolor, 132, 133, 137, var. **bracteata**, 134, 137.  
*Secale cereale*, 221.  
*Selaginella*, 68; **apoda**, 68; apus, 68.  
*Sedum roseum*, 154.  
*Senecio Robbinsii*, 154; *sylvaticus*, 148; *vulgaris*, 36, 37.  
*Setaria glauca*, 212; *viridis*, 226.  
*Seymeria*, 134; **cassioides**, 134, 137; *tenuifolia*, 134, 137.  
 Shields, M. A., The field Meeting of the Vermont Botanical and Bird Clubs, 143.  
 Shrubs, A handy Book on Trees and, 103.  
*Silene antirrhina*, 96, 97, Two Variations of, 96, var. **confinis**, 97, forma **Deaneana**, 96, var. *depauperata*, 97, var. *divaricata*, 97, var. *laevigata*, 97, var. *vaccariifolia*, 97.  
 Silver Fir, 165.  
*Sisymbrium*, 138; *altissimum*, 30, 35; *brachycarpon*, 138-140, and Allies, 138, var. **filipes**, 140, 141, var. **intermedium**, 140, 141; *canescens*, 138-140, var. *brachycarpon*, 138, 140; *incisum*, 30, 139, 140, var. *filipes*, 139, 141; *officinale*, var. *leiocarpum*, 35; *pinnatum*, 140; *Sophia*, 140.  
*Sisyrinchium angustifolium*, 153.  
*Sium Berula*, 131; *cicutaeifolium*, 131, 137, 150, 217; *suave*, 131, 132, 137.  
*Smilacina trifolia*, 149.  
*Smilax Bona-nox*, 64; *herbacea*, 63; *pulverulenta*, 63.  
 Smith, C. P., *Carex Tuckermani niagarensis*; a neglected Sedge, 57.  
 Society of Maine, The Josselyn Botanical, 128; of Maine, The twenty-first annual Meeting of the Josselyn Botanical, 203.  
*Solanum Dulcamara*, 217; *rostratum*, 30.  
*Solidago altissima*, 10, 12, 218; *angustifolia*, 27; *aspera*, 7; *azorica*, 27; **Bartramiana**, 10, 11; *bicolor*, 36; *caesia*, 36; *canadensis*, 8-12, 36, 218, var. *gilvocanescens*, 9, 12, var. **Hargerii**, 11; **chlorolepis**, 3; *confinis*, 27; *decumbens*, 4; *Elliottii*, 7, var. *divaricata*, 7; *elongata*, 8, 9; *erecta*, 2; *floribunda*, 12; *graminifolia*, 12, 151, 218, var. **camporum**, 12, var. *Nuttallii*, 13, var. **polycephala**, 12, var. **septentrionalis**, 12; *Guiradonis*, 5; *hispida*, 1, var. *arnoglossa*, 2, 3, var. **disjuncta**, 2, var. *fallax*, 3, var. *lanata*, 1, var. **tonsa**, 2; *humilis*, 6, var. **peracuta**, 6; *humilius*, 6; in the White Mountains, A northern, 72; *juncea*, 36; **Klughii**, 5; *lepida*, 8-11, var. **elongata**, 9, 10, var. *fallax*, 9, 11, var. **molina**, 9; *macrophylla*, 152, 153; **mensalis**, 4; *multi-radiata*, 4, 5, var. **arctica**, 4; *nemorialis*, 36; *polycephala*, 12; *racemosa*, 6, 204; *Randii*, 148; *rugosa*, 7, 12, 36, 152, 154, var. **aspera**, 7, 218, var. *villosa*, 7, 8, 72; *sempervirens*, 26, 27, 36, 147, 208, 218, 221, An insular Variety of, 26, var. **azorica**, 27; *serotina*, 8-10, var. *salebrosa*, 9; *stricta*, 26, var. *angustifolia*, 26; *uliginosa*, 6; *uniligulata*, 6, 7, 149, var. **levipes**, 7; *Virgaurea*,  $\mu$  *arctica*, 4.  
 Some Algae from the Chinha Islands, 89; anomalous Species and Varieties of *Bidens* in eastern North America, 20; neglected Names in Walter's Flora Caroliniana, 129; new or unrecorded Compositae chiefly of northeastern America, 1.  
*Sonchus arvensis*, 220; *asper*, 36; *oleraceus*, 36, 220.  
*Sophia brachycarpa*, 138; *canescens*, 138; *intermedia*, 139-141; *magna*, 138; *pinnata*, 138.  
 Sorrel, 226.  
 Spanish Chestnuts, 49.  
*Sparganium angustifolium*, 154; *fluctuans*, 150; *minimum*, 65.  
*Spartina*, 212; *glabra*, var. *alterniflora*, 147, 209, 212, 221, var. *pilosa*, 212, 221; *Michauxiana*, 146, 212; *patens*, 38, 147, 212, 214, 221, var. *juncea*, 213, 221.  
*Specularia*, 233; *perfoliata*, 30, 233.  
*Spergula arvensis*, 35.  
*Spergularia canadensis*, 147; *rubra*, 35, 227; *salina*, 215, 221.  
*Sphaerocarpos*, 109.  
*Spinacea*, 64, 178; *oleracea*, 178.  
*Spiraea latifolia*, 150, 153; *tomentosa*, 143, 153, A white-flowered, 142, forma **albiflora**, 143; *trifoliata*, 130.  
*Spiranthes cernua*, 34; *Romanzofiana*, 153, 154.

- Sporobolus asperifolius*, 160; *Richardsonis*, 204.  
*Spruce*, 65; *Hemlock*, 59-61, 164-168, The Name of the, 59; *White*, 61, 167, 168.  
*Stellaria borealis*, 151; *graminea*, 151; *humifusa*, 147; *media*, 35.  
*Streblocladia camptoclada*, 95; *spicata*, 95.  
*Streptopus amplexifolius*, 154; *roseus*, 152.  
*Struthiopteris*, 161; *germanica*, 161, 162, var. *pennsylvanica*, 162-164; *nodulosa*, 162, 164; *pennsylvanica*, 162, 164.  
*Suaeda*, 178; *linearis*, 178, 215, 221; *maritima*, 84, 146, 147, 178; *Richii*, 178.  
*Subularia*, 155.  
*Swertia difformis*, 50, 51.  
*Syntherisma*, 130; *villosa*, 130.  
*Syringa*, Common, 121.  
Systematic Studies on *Oenothera*,—V, 41.  
*Tanacetum vulgare*, 36, 219.  
*Taraxacum*, 227; *officinale*, 36, 209, 220.  
*Taxus canadensis*, 152.  
Taylor, N. [Notice of Work], 62.  
Tea, Labrador, 149.  
Teratological specimen of *Panicum amarulum* Hitchc. & Chase, 72.  
*Thalictrum polygamum*, 151, var. *hebecarpum*, 151.  
*Thlaspi arvense*, 30.  
Three of Clayton's Oaks in the British Museum, 39.  
*Thuya occidentalis*, 152.  
*Thymra caroliniana*, 132.  
*Thysanthema semiflosculare*, 136.  
*Tofieldia glutinosa*, 204.  
Torrey, G. S., The Varieties of *Cardamine oligosperma*, 156.  
Trees and Shrubs, a handy Book on, 103.  
*Tridens*, 72.  
*Trientalis americana*, 152.  
*Trifolium*, 32; *arvense*, 38; *echinatum*, 32; *hybridum*, 35, 216; *pratense*, 35, 216; *repens*, 35.  
*Triglochin maritima*, 147, 208, 211, 221; *palustris*, 64.  
*Trillium undulatum*, 152.  
*Trisetum spicatum*, 65.  
*Tsuga americana*, 59, 168; *canadensis*, 59, 60; *caroliniana*, 165.  
*Tubiflora caroliniensis angustifolia*, 131.  
Twentieth Anniversary of the New York Botanical Garden, 144.  
Twenty-first annual Meeting of the Josselyn Botanical Society of Maine, 203.  
Two Extensions of Range in Gramineae, 71; more foreign Plants found on Wool-waste at Westford, 32; *Philadelphus Combinations*, 121; Variations of *Silene antirrhina*, 96.  
*Typha latifolia*, 150, 209, 210.  
*Ulmus*, 170; *americana*, 170; *campestris*, 170; *fulva*, 170; *scabra*, 170.  
*Ulva*, 91.  
*Urtica*, 171; *dioica*, 171; *gracilis*, 171; *Lyallii*, 64, 171; *urens*, 171.  
Urticaceae, 169.  
*Vaccinium canadense*, 153; *macrocarpon*, 148; *Oxycoccus*, 148; *pennsylvanicum*, 153, var. *nigrum*, 153; *Vitis-Idaea*, var. *minus*, 148.  
*Valeriana*, 66; *uliginosa*, 66.  
Variations of *Silene antirrhina*, Two, 96.  
Varieties of *Cardamine oligosperma*, 156.  
*Verbena hastata*, 217.  
Vermont Botanical and Bird Clubs, The field Meeting of the, 143; Botanical Club, The 20th annual winter Meeting of the, 86.  
*Vernonia acaulis*, 135, 137; *oligophylla*, 135, 137.  
*Veronica officinalis*, 152; *peregrina*, 218.  
*Viburnum cassinoides*, 152; *Lentago*, 152.  
*Vicia angustifolia*, var. *segetalis*, 148.  
*Vincetoxicum*, 130.  
*Viola*, 71, 202; *adunca*, 71; *cucullata*, 70, 151, forma *albiflora*, 180; *cuspidata*, 70; *incognita*, 151, 152, 154; *labradorica*, 154; *lanceolata*, 151, 217; *nephrophylla*, 71; *pallens*, 151, 152; *palustris*, 71; *pedata*, 35, 37, forma *alba*, 180; *primulifolia*, 151, 154; *renifolia*, 71; *rostrata*, 180, An *albino*, 180, forma **Phelpsiae**, 180; *Selkirkii*, 71; *septentrionalis*, 70, 154, in British Columbia, 70; *sororia*, 70; *tricolor*, 35, 38.  
Walter's Flora Caroliniana, Some neglected Names in, 129.



- Washington and Michaux, 49.  
Weatherby, C. A., Two Extensions of Range in Gramineae, 71.  
Webster, H., A rash Mycophagist, 30.  
Weed Growth and unusual Rainfall, 224.  
Weeds growing in Amherst, 29.  
Westford, Two more foreign Plants found on Wool-waste at, 32.  
White flowered *Spiraea tomentosa*, 142; Mountains, *Arenaria stricta* in the, 232; Mountains, A northern *Solidago* in the, 72; Spruce, 61, 167, 168.  
Wiegand, K. M., The Genus *Euphrasia* in North America, 181.  
Wilcox, A. W., The field Meeting of the Vermont Botanical and Bird Clubs, 143.  
Witch Grass, 225.  
Woodward, R. W., *Bidens connata* petiolata, 234.  
*Woodwardia areolata*, 162.  
Wool-waste at Westford, Two more foreign Plants found on, 32.  
*Xanthium echinatum*, 219.  
*Xyris montana*, 65.  
*Zizania palustris*, 212.  
*Zostera marina*, 147.









## IMPORTANT NEW BOTANICAL BOOKS

COULTER'S PLANT LIFE AND PLANT USES . . . . .	\$1.20
By John Gaylord Coulter, Ph. D., Critic Teacher of Biology, University High School, Normal, Illinois.	
Edition with Cowles and Coulter's Spring Flora . . . . .	1.50
COULTER'S BOTANICAL NOTEBOOK AND LABORA- TORY MANUAL . . . . .	.36
By John Gaylord Coulter, Ph. D.	
With loose leaf binder . . . . .	.72
COWLES AND COULTER'S SPRING FLORA . . . . .	.60
By Henry C. Cowles, Ph. D., Associate Professor of Plant Ecology, University of Chicago, and John Gaylord Coulter, Ph. D.	
Edition with Andrews's Practical Botany . . . . .	1.50

DESCRIPTIVE CIRCULARS SENT ON REQUEST

AMERICAN BOOK COMPANY

New York

Cincinnati

Chicago

Boston

## HOME GROWN LILIES.

Fresh from Beds. Wild Flowers, Hardy plants of all kinds.

Send for catalogue

F. H. HORSFORD, Charlotte, Vt.

**A YOSEMITE FLORA** by Harvey Monroe Hall and Car-  
lotta Case Hall. A descriptive guide,  
with keys. Includes most species of the Sierra Nevada Mountains.  
282 pages, 11 plates, and 170 text-figures. Pocket-size. Attractively  
bound in flexible leather. Postpaid, \$2.00 net.

PAUL ELDER & CO.

239 GRANT AVENUE, SAN FRANCISCO, CAL.

Mounting Papers, Driers, Genus Covers, Collecting  
Cases, Plant Presses

BOTANICAL CATALOG 91. DO YOU HAVE IT?

CAMBRIDGE BOTANICAL SUPPLY COMPANY

Laboratory . . . . . WAVERLEY, MASS. . . . . Equipment  
(Mention Rhodora)